BEST AVAILABLE COPY



IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Baker et al.

Docket No:

39780-2830C1P10

Serial No:

10/006,768

Group Art Unit:

1647

Filed:

December 6, 2001

Examiner:

Rachel B. Kapust

For:

SECRETED AND TRANSMEMBRANE POLYPEPTIDES AND NUCLEIC

ACIDS ENCODING THE SAME

Commissioner for Patents Washington, D.C. 20231

DECLARATION OF LUC DESNOYERS, Ph.D., DR. AUDREY GODDARD, Ph.D., DR. PAUL J. GODOWSKI, Ph.D., DR. AUSTIN GURNEY, Ph.D., DR. COLIN K. WATANABE and DR. WILLIAM WOOD, Ph.D. UNDER 37 CFR 1.131

- 1. We are the inventors of the above-identified application.
- 2. We have read and understood the claims pending in this application, and are aware that the claims have been rejected as anticipated by International Patent Application Publication No. WO 00/00610 (Lal *et al.*, publication date January 6, 2000).
- 3. We conceived and reduced to practice the invention claimed in the above-identified application in the United States prior to January 6, 2000.
- 4. At the time the present invention was made, one of the inventors, Luc Desnoyers, Ph.D., was, as still is, responsible for overseeing the testing of novel polypeptides, including the polypeptide designated PRO1412, in chondrocyte proliferation assay (Assay #111, Example 153). This assay is used to find agents that are capable of inducing chondrocyte proliferation and/or redifferentiation, and can, therefore, be used in the treatment of joint diseases using a tissue engineering approach or as promising drug candidates to repair aging or arthritic joints, for example, in which the chondrocytes have been dedifferentiated.

- In this assay, isolated chondrocyte cells are seeded in 96 well plates with either serum-free medium (negative control), staurosporin (positive control) or the test PRO polypeptide. After 5 days, fluorescence dye is added to each plate and measured. The readout of the fluorescence from a plate containing the serum-free medium is measured to establish a background fluorescence level. A positive result in the assay is obtained when the fluorescence of the PRO polypeptide-treated sample is more like that of the positive control than the negative control. This type of fluorescence determination, wherein the readout is compared to positive and negative controls, is well known in the art.
- 6. A copy of a page from the Genengenes database which reports a positive result for the PRO1412 polypeptide encoded by DNA 64897-1628 (UNQ730) in Assay #111 is attached to this declaration (with its date redacted) as Exhibit A. The positive results reported in the database were also obtained prior to January 6, 2000.
- 7. Copies of pages from laboratory notebook showing the positive results for the PRO1412 polypeptide (SEQ ID NO:140), identified by Pin number PIN753-1, in Assay #111 are attached to this declaration (with dates redacted) as Exhibit B. These experiments were performed and the results were obtained prior to January 6, 2000.
- 8. Exhibits A and B clearly show that the polypeptide designated PRO1412 was tested, and its ability to induce the proliferation and/or redifferentiation was determined prior to January 6, 2000.

Luc Desnoyers, Ph.D.	08/14/2004 Date
Audrey Goddard, Ph.D.	Date
Paul J. Godowski, Ph.D.	Date
Austin Gurney, Ph.D.	Date
Colin K. Watanabe	Date
William Wood, Ph.D.	 Date

SV 2055606 v1 8/18/04 3:22 PM (39780.2830)



Baker et al.

Docket No:

39780-2830C1P10

Serial No:

10/006,768

Group Art Unit:

1647

Filed:

December 6, 2001

Examiner:

Rachel B. Kapust

For:

SECRETED AND TRANSMEMBRANE POLYPEPTIDES AND NUCLEIC

ACIDS ENCODING THE SAME

Commissioner for Patents Washington, D.C. 20231

DECLARATION OF LUC DESNOYERS, Ph.D., DR. AUDREY GODDARD, Ph.D., DR. PAUL J. GODOWSKI, Ph.D., DR. AUSTIN GURNEY, Ph.D., DR. COLIN K. WATANABE and DR. WILLIAM WOOD, Ph.D. **UNDER 37 CFR 1.131**

- 1. We are the inventors of the above-identified application.
- 2. We have read and understood the claims pending in this application, and are aware that the claims have been rejected as anticipated by International Patent Application Publication No. WO 00/00610 (Lal et al., publication date January 6, 2000).
- 3. We conceived and reduced to practice the invention claimed in the above-identified application in the United States prior to January 6, 2000.
- 4. At the time the present invention was made, one of the inventors, Luc Desnoyers, Ph.D., was, as still is, responsible for overseeing the testing of novel polypeptides, including the polypeptide designated PRO1412, in chondrocyte proliferation assay (Assay #111, Example 153). This assay is used to find agents that are capable of inducing chondrocyte proliferation and/or redifferentiation, and can, therefore, be used in the treatment of joint diseases using a tissue engineering approach or as promising drug candidates to repair aging or arthritic joints, for example, in which the chondrocytes have been dedifferentiated.

- 5. In this assay, isolated chondrocyte cells are seeded in 96 well plates with either serum-free medium (negative control), staurosporin (positive control) or the test PRO polypeptide. After 5 days, fluorescence dye is added to each plate and measured. The readout of the fluorescence from a plate containing the serum-free medium is measured to establish a background fluorescence level. A positive result in the assay is obtained when the fluorescence of the PRO polypeptide-treated sample is more like that of the positive control than the negative control. This type of fluorescence determination, wherein the readout is compared to positive and negative controls, is well known in the art.
- 6. A copy of a page from the Genengenes database which reports a positive result for the PRO1412 polypeptide encoded by DNA 64897-1628 (UNQ730) in Assay #111 is attached to this declaration (with its date redacted) as Exhibit A. The positive results reported in the database were also obtained prior to January 6, 2000.
- 7. Copies of pages from laboratory notebook showing the positive results for the PRO1412 polypeptide (SEQ ID NO:140), identified by Pin number PIN753-1, in Assay #111 are attached to this declaration (with dates redacted) as Exhibit B. These experiments were performed and the results were obtained prior to January 6, 2000.
- 8. Exhibits A and B clearly show that the polypeptide designated PRO1412 was tested, and its ability to induce the proliferation and/or redifferentiation was determined prior to January 6, 2000.

Luc Desnoyers, Ph.D.	Date
Q- Stoddard	8/19/04
Audrey Goddard, Ph.D.	Date
r.	
Paul J. Godowski, Ph.D.	Date
Austin Gurney, Ph.D.	Date
Colin K. Watanabe, Ph.D.	Date
	_·
William Wood, Ph.D.	Date

SV 2055606 v1 8/18/04 2:52 PM (39780.2830)



Baker et al.

Docket No:

39780-2830C1P10

Serial No:

10/006,768

Group Art Unit:

1647

Filed:

December 6, 2001

Examiner:

Rachel B. Kapust

For:

SECRETED AND TRANSMEMBRANE POLYPEPTIDES AND NUCLEIC

ACIDS ENCODING THE SAME

Commissioner for Patents Washington, D.C. 20231

DECLARATION OF LUC DESNOYERS, Ph.D., DR. AUDREY GODDARD, Ph.D., DR. PAUL J. GODOWSKI, Ph.D., DR. AUSTIN GURNEY, Ph.D., DR. COLIN K. WATANABE and DR. WILLIAM WOOD, Ph.D. UNDER 37 CFR 1.131

- 1. We are the inventors of the above-identified application.
- 2. We have read and understood the claims pending in this application, and are aware that the claims have been rejected as anticipated by International Patent Application Publication No. WO 00/00610 (Lal *et al.*, publication date January 6, 2000).
- 3. We conceived and reduced to practice the invention claimed in the above-identified application in the United States prior to January 6, 2000.
- 4. At the time the present invention was made, one of the inventors, Luc Desnoyers, Ph.D., was, as still is, responsible for overseeing the testing of novel polypeptides, including the polypeptide designated PRO1412, in chondrocyte proliferation assay (Assay #111, Example 153). This assay is used to find agents that are capable of inducing chondrocyte proliferation and/or redifferentiation, and can, therefore, be used in the treatment of joint diseases using a tissue engineering approach or as promising drug candidates to repair aging or arthritic joints, for example, in which the chondrocytes have been dedifferentiated.

- In this assay, isolated chondrocyte cells are seeded in 96 well plates with either serum-free medium (negative control), staurosporin (positive control) or the test PRO polypeptide. After 5 days, fluorescence dye is added to each plate and measured. The readout of the fluorescence from a plate containing the serum-free medium is measured to establish a background fluorescence level. A positive result in the assay is obtained when the fluorescence of the PRO polypeptide-treated sample is more like that of the positive control than the negative control. This type of fluorescence determination, wherein the readout is compared to positive and negative controls, is well known in the art.
- 6. A copy of a page from the Genengenes database which reports a positive result for the PRO1412 polypeptide encoded by DNA 64897-1628 (UNQ730) in Assay #111 is attached to this declaration (with its date redacted) as Exhibit A. The positive results reported in the database were also obtained prior to January 6, 2000.
- 7. Copies of pages from laboratory notebook showing the positive results for the PRO1412 polypeptide (SEQ ID NO:140), identified by Pin number PIN753-1, in Assay #111 are attached to this declaration (with dates redacted) as Exhibit B. These experiments were performed and the results were obtained prior to January 6, 2000.
- 8. Exhibits A and B clearly show that the polypeptide designated PRO1412 was tested, and its ability to induce the proliferation and/or redifferentiation was determined prior to January 6, 2000.

Luc Desnoyers, Ph.D.	Date
Audrey Goddard, Ph.D. Paul J. Godowski, Ph.D.	Date S/19/0'/ Date
Austin Gurney, Ph.D.	Date
Colin K. Watanabe	Date
William Wood, Ph.D.	Date

SV 2055606 v1 8/18/04 3:22 PM (39780.2830)



Baker et al.

Docket No:

39780-2830C1P10

Serial No:

10/006,768

Group Art Unit:

1647

Filed:

December 6, 2001

Examiner:

Rachel B. Kapust

For:

SECRETED AND TRANSMEMBRANE POLYPEPTIDES AND NUCLEIC

ACIDS ENCODING THE SAME

Commissioner for Patents Washington, D.C. 20231

DECLARATION OF LUC DESNOYERS, Ph.D., DR. AUDREY GODDARD, Ph.D., DR. PAUL J. GODOWSKI, Ph.D., DR. AUSTIN GURNEY, Ph.D., DR. COLIN K. WATANABE and DR. WILLIAM WOOD, Ph.D. UNDER 37 CFR 1.131

- 1. We are the inventors of the above-identified application.
- 2. We have read and understood the claims pending in this application, and are aware that the claims have been rejected as anticipated by International Patent Application Publication No. WO 00/00610 (Lal *et al.*, publication date January 6, 2000).
- 3. We conceived and reduced to practice the invention claimed in the above-identified application in the United States prior to January 6, 2000.
- 4. At the time the present invention was made, one of the inventors, Luc Desnoyers, Ph.D., was, as still is, responsible for overseeing the testing of novel polypeptides, including the polypeptide designated PRO1412, in chondrocyte proliferation assay (Assay #111, Example 153). This assay is used to find agents that are capable of inducing chondrocyte proliferation and/or redifferentiation, and can, therefore, be used in the treatment of joint diseases using a tissue engineering approach or as promising drug candidates to repair aging or arthritic joints, for example, in which the chondrocytes have been dedifferentiated.

- 5. In this assay, isolated chondrocyte cells are seeded in 96 well plates with either serum-free medium (negative control), staurosporin (positive control) or the test PRO polypeptide. After 5 days, fluorescence dye is added to each plate and measured. The readout of the fluorescence from a plate containing the serum-free medium is measured to establish a background fluorescence level. A positive result in the assay is obtained when the fluorescence of the PRO polypeptide-treated sample is more like that of the positive control than the negative control. This type of fluorescence determination, wherein the readout is compared to positive and negative controls, is well known in the art.
- 6. A copy of a page from the Genengenes database which reports a positive result for the PRO1412 polypeptide encoded by DNA 64897-1628 (UNQ730) in Assay #111 is attached to this declaration (with its date redacted) as Exhibit A. The positive results reported in the database were also obtained prior to January 6, 2000.
- 7. Copies of pages from laboratory notebook showing the positive results for the PRO1412 polypeptide (SEQ ID NO:140), identified by Pin number PIN753-1, in Assay #111 are attached to this declaration (with dates redacted) as Exhibit B. These experiments were performed and the results were obtained prior to January 6, 2000.
- 8. Exhibits A and B clearly show that the polypeptide designated PRO1412 was tested, and its ability to induce the proliferation and/or redifferentiation was determined prior to January 6, 2000.

Luc Desnoyers, Ph.D.	Date
Audrey Goddard, Ph.D.	Date
Paul J. Godowski, Ph.D. Austin Gurney, Ph.D.	Date S/70/59 Date
Colin K. Watanabe, Ph.D.	Date
William Wood, Ph.D.	Date

SV 2055606 v1 8/18/04 2:52 PM (39780.2830)



Baker et al.

Docket No:

39780-2830C1P10

Serial No:

10/006,768

Group Art Unit:

1647

Filed:

December 6, 2001

Examiner:

Rachel B. Kapust

For:

SECRETED AND TRANSMEMBRANE POLYPEPTIDES AND NUCLEIC

ACIDS ENCODING THE SAME

Commissioner for Patents Washington, D.C. 20231

DECLARATION OF LUC DESNOYERS, Ph.D., DR. AUDREY GODDARD, Ph.D., DR. PAUL J. GODOWSKI, Ph.D., DR. AUSTIN GURNEY, Ph.D., DR. COLIN K. WATANABE and DR. WILLIAM WOOD, Ph.D. UNDER 37 CFR 1.131

- 1. We are the inventors of the above-identified application.
- 2. We have read and understood the claims pending in this application, and are aware that the claims have been rejected as anticipated by International Patent Application Publication No. WO 00/00610 (Lal *et al.*, publication date January 6, 2000).
- 3. We conceived and reduced to practice the invention claimed in the above-identified application in the United States prior to January 6, 2000.
- 4. At the time the present invention was made, one of the inventors, Luc Desnoyers, Ph.D., was, as still is, responsible for overseeing the testing of novel polypeptides, including the polypeptide designated PRO1412, in chondrocyte proliferation assay (Assay #111, Example 153). This assay is used to find agents that are capable of inducing chondrocyte proliferation and/or redifferentiation, and can, therefore, be used in the treatment of joint diseases using a tissue engineering approach or as promising drug candidates to repair aging or arthritic joints, for example, in which the chondrocytes have been dedifferentiated.

- 5. In this assay, isolated chondrocyte cells are seeded in 96 well plates with either serum-free medium (negative control), staurosporin (positive control) or the test PRO polypeptide. After 5 days, fluorescence dye is added to each plate and measured. The readout of the fluorescence from a plate containing the serum-free medium is measured to establish a background fluorescence level. A positive result in the assay is obtained when the fluorescence of the PRO polypeptide-treated sample is more like that of the positive control than the negative control. This type of fluorescence determination, wherein the readout is compared to positive and negative controls, is well known in the art.
- 6. A copy of a page from the Genengenes database which reports a positive result for the PRO1412 polypeptide encoded by DNA 64897-1628 (UNQ730) in Assay #111 is attached to this declaration (with its date redacted) as Exhibit A. The positive results reported in the database were also obtained prior to January 6, 2000.
- 7. Copies of pages from laboratory notebook showing the positive results for the PRO1412 polypeptide (SEQ ID NO:140), identified by Pin number PIN753-1, in Assay #111 are attached to this declaration (with dates redacted) as Exhibit B. These experiments were performed and the results were obtained prior to January 6, 2000.
- 8. Exhibits A and B clearly show that the polypeptide designated PRO1412 was tested, and its ability to induce the proliferation and/or redifferentiation was determined prior to January 6, 2000.

Luc Desnoyers, Ph.D.	Date
Audrey Goddard, Ph.D.	Date
Paul J. Godowski, Ph.D.	Date
Austin Gurney, Ph.D.	Date Ang 24 not
Colin K. Watanabe, Ph.D.	Date
William Wood Ph D	— Date

SV 2055606 v1 8/18/04 2:52 PM (39780.2830)



Applicants:

Baker et al.

Docket No:

39780-2830C1P10

Serial No:

10/006,768

Group Art Unit:

1647

Filed:

December 6, 2001

Examiner:

Rachel B. Kapust

For:

SECRETED AND TRANSMEMBRANE POLYPEPTIDES AND NUCLEIC

ACIDS ENCODING THE SAME

Commissioner for Patents Washington, D.C. 20231


DECLARATION OF LUC DESNOYERS, Ph.D., DR. AUDREY GODDARD, Ph.D., DR. PAUL J. GODOWSKI, Ph.D., DR. AUSTIN GURNEY, Ph.D., DR. COLIN K. WATANABE and DR. WILLIAM WOOD, Ph.D. **UNDER 37 CFR 1.131**

- 1. We are the inventors of the above-identified application.
- We have read and understood the claims pending in this application, and are aware 2: that the claims have been rejected as anticipated by International Patent Application Publication No. WO 00/00610 (Lal et al., publication date January 6, 2000).
- 3. We conceived and reduced to practice the invention claimed in the above-identified application in the United States prior to January 6, 2000.
- 4. At the time the present invention was made, one of the inventors, Luc Desnoyers, Ph.D., was, as still is, responsible for overseeing the testing of novel polypeptides, including the polypeptide designated PRO1412, in chondrocyte proliferation assay (Assay #111, Example 153). This assay is used to find agents that are capable of inducing chondrocyte proliferation and/or redifferentiation, and can, therefore, be used in the treatment of joint diseases using a tissue engineering approach or as promising drug candidates to repair aging or arthritic joints, for example, in which the chondrocytes have been dedifferentiated.

- In this assay, isolated chondrocyte cells are seeded in 96 well plates with either serum-free medium (negative control), staurosporin (positive control) or the test PRO polypeptide. After 5 days, fluorescence dye is added to each plate and measured. The readout of the fluorescence from a plate containing the serum-free medium is measured to establish a background fluorescence level. A positive result in the assay is obtained when the fluorescence of the PRO polypeptide-treated sample is more like that of the positive control than the negative control. This type of fluorescence determination, wherein the readout is compared to positive and negative controls, is well known in the art.
- 6. A copy of a page from the Genengenes database which reports a positive result for the PRO1412 polypeptide encoded by DNA 64897-1628 (UNQ730) in Assay #111 is attached to this declaration (with its date redacted) as Exhibit A. The positive results reported in the database were also obtained prior to January 6, 2000.
- 7. Copies of pages from laboratory notebook showing the positive results for the PRO1412 polypeptide (SEQ ID NO:140), identified by Pin number PIN753-1, in Assay #111 are attached to this declaration (with dates redacted) as Exhibit B. These experiments were performed and the results were obtained prior to January 6, 2000.
- 8. Exhibits A and B clearly show that the polypeptide designated PRO1412 was tested, and its ability to induce the proliferation and/or redifferentiation was determined prior to January 6, 2000.

Luc Desnoyers, Ph.D.	Date
Audrey Goddard, Ph.D.	Date
Paul J. Godowski, Ph.D.	Date
Austin Gurney, Ph.D.	Date
Colin K. Watanabe	Date
William Wood, Ph.D.	Date

SV 2055606 v1 8/18/04 3:22 PM (39780.2830)



NSY I DINA I DOM I EXEL FEMILIE, SI LIBI I LOTI I MARI I QUI I PRIBI I PRO I PUBLI I SACI UNO I XELLI MARI I SACI MARI I SACI UNO I XELLI MARI I SACI MARI I SACI

GenenGenes Feedback


Primary Assay Result Assay ID Assay Name Assay Oate

Notebook Num	XXXXXX						1.00	1%	1%	1%	1%	1%
	1%	1%	1%	1%	1%	1%	170	100		10	11.	12
	1	7 2	3	4	5	6	1 7		7	PIN742-1	PIN746-1	PIN750-1
	Staurosponn	Staurosporin	Staurosportn	PIN717-1	PIN721-1	PIN725-1	PIN730-1	PIN734-1	PIN738-1	PINTAZ-I		
Ĩ.	Media	Media	Media					PIN735-1	PIN739-1	PIN743-1	PIN747-1	PIN751-1
· č	PIN708-1	PIN711-1	PIN714-1	PIN716-1	PIN722-1	PIN727-1	PIN731-1	PIR/ 33-1	7-447-43-1	7		
0		i					PIN732-1	PIN736-1	P1N740-1	PIN744-1	PIN748-1	PIN752-1
· E	PIN709-1	PIN712-1 -	PIN715-1	PIN719-1	PIN724-1	PIN728-1	P107.32-1	F 1141 20-1				
							PIN733-1	P1N737-1	PIN741-1	PIN745-1	PtN749-1	PIN753-1
o ·	PIN710-1	PIN713-1	PIN715-1	PIN720-1	PIN725-1	PIN729-1	JE1171 33-1	,	1	1	L	<u> </u>
. н	1											

	• •							7			10	1 11	12	į .
			2	3	4				103.1	74.4	0.58	155.6	82.1	i.
		44.1	57.1	98.4	100,4	173.2	166.5	166.6	104.1	14.4	104.1	78.5	119.8	Ĺ.
	l ^ L			94.4	144.4	98.6	103.6	118.8	75.5	89.3				ĺ
-	1 'B L	81.1	·159.7	54.4		68.5	64.9	60.4	56.6	67.8	50.6	63.7	107.2	
- 1	1 e · . [. 85.4	91.8	89.5	69.1	60.3		60.3	56.4	70.9	75.3	79.5	119.6	
		84.8	102.4	_75.6	.704	54.7	3/			· 74.3	72.1	57.5	175.3	ł
	r 2	72.4	73.5	58.0	71.4	55.9	51.1	59.7	47.5		1	64.5	180.5	ı.
		102.9		69.7	131.0	84.7	83.8	75.2	55.8	58.6				1
	1. F 1	59.9	102.3				57.6	64,6	73.5	77.2	64.9	182.5	183.7	1.4
		116.0	125.9	159.3	105.5	77.5		128.4	125.7	95.4	137.9	172.1	. 139.5	
	1 2 1	. 144.3	102.0	192.0	193.6	181.4	124.0	1 120.4						-

Conc			1.00%	ļ			
PIN	N1	N2	AVERAGE	STOEV .	Positive	Verified	Comments
PINTOA-1	0,788	0.780	0.784	0.0			
PIN709-1	0.949	0.645	0.797	0.2	1 .	1	1
PIN710-1	1,070	1.331	1.201	0.2			
PIN711-1	0.847	0.945	0.896	0.1	1	1	
PIN712-1	0.578	0.944	Q.811'	0.2	1 :		1
PIN713-1	1,162	0.941	1.052	0.2			1
PIN714-1	0.825	0.697	0.782	0.1	1 1	* ** .	
	0.535	0.744	0.640	0.1			
PIN715-1	. 1,469	1,771	1,620	0.2	Pasitive	1	L '
PIN716-1	0.925	1,333	1,129	0.3	4	1	
PIN717-1	0.622	0.653	0.738	0.1			
PIN718-1		1,216	0.938	0.4			
PIN719-1	.0.839	1.786	1,380	0.0	Positive		
PIN720-1	0.973	0.910	1.254	0.5		1	1 '
PIN721-1	1.596		0.614	0.0	1.		4 3 5
PIN722-1	0.632	0.597	0.648	0.2	1		1
PIN724-1	0.515	0.781	1.102	0.5	1 .	T	4
PIN725-1	0.715	1,459		0.4	.1 : :		1 .
PIN726-1	. 1,537	0.958	1.246			1	
PIN727-1	0,599	0.343	0.471	0.2	1 .	· .	
PIN728-1	0.471	0.774	0.523	0.2			1
PIN729-1	0.532	1,144	. 0.838	0.4			1 .
PIN730-1	1.538	1,096	1,317	0.3	Positive		4 .
PIN731-1	0.557	0.556	0.557				
PIN732-1	0.551	0,722	. 0.636	0.1		1	ļ .
PIN733-1	. 0.595	1,184	0.890	0.4		I .	
PIN734-1	0.951	0,697	0.824	0.2	· · ·	1	
PIN735-1	0.522	0,520	0.521	0.0	ļ.	l .	1
PIN736-1	0.438	0,617	0.527	0.1	1 '	1	1
	0,678	1,159	0.919	0.3	. i .		1
PIN737-1	0.586	0.624	0.755	. 0.1	1 .	1	1-
PIN738-1	0.524	0.654	0.639	0.0	1 .	1	
P!N739-1	0.688	0.635	0.660	. 0.0	- A	1	1
PIN740-1	0.712	0.880	0.796	0.1	` l .	· .	
PIN741-1		0.961	0.886	0.1	i i	1 .	. [
PIN742-1	0.812	0.695	0.518	0.1	1	١.	1
PIN743-1	0.541	0.751	0.708	0.1	. ' ''		.)
PIN744-1	0.665		0.935	0.5	- 1°		Ι.
PIN745-1	0.599	1,272	1.020	0.5		1	1
PIN746-1	1,436	0.724		0.1		- I	1 4 - 5
PIN747-1	0.588	0.733	0.661	0.2		F.	
PIN748-1	0.464	0.781		0.2	Positive	1	1
PIN749-1	1.584	1,580	1,636		LOSIGNA	.: .	١.
PIN750-1	0.757	1.105 .	0,931	0.2		3.1	1 1
PIN751-1	0.989	1,104	1.048	0.1		11 · .	4
PIN752-1	1.518	1.565	1.842	. 0.0	Positive	1 .	.11
PIN753-1	1.695	1.287	1,491	0.3	Positive	_ .	

PINE		Average	STOEV
PIN708-1		0.78	0.0
PIN709-1 .		0.60	0.2 .
PIN710-1		1.20	0.2
PIN711-1		0.90	0.1
PIN712-1	. 1	0.81	0.2
PIN713-1		1.05	.0.2
PIN714-1	1	. 0,75	·*O.1
PIN7 15-1		0.64	· 0.1
PIN716-1		1.62	0.2
PIN717-1		1,13	0.3
PIN716-1		0.74	0.1
PIN719-1		0.94	0.4
1N720-1	1	1,38	0.6
PIN721-1		1.25	0.5
1N722-1		0.61	0.0
7N724-1 7N725-1		0.65	0.2
71N725-1 71N726-1		1.10	0.5
1N725-1		1.25	0.4
7N/2/-1 PIN728-1	. 1	0.47	0.2
1N728-1 PIN729-1		0.62	0.2
PIN730-1		0.84	0.4
*1N730-1 *1N731-1 ·	· . i	1.32	0.3
7N/31-1 ·		. 0.56 0.64	0.0
2N732-1		0.64	0.1
7N734-1		0.89	0.4
1N735-1		0.52	0.2
1N738-1	· i	0.52	0.1
1N737-1		0.92	0.3
PIN738-1		0.76	0.1
1N739-1	. 1	0.64	0.0
1N740-1		0.66	0.0
PIN741-1		0.80	0.1
MN742-1		0.89	0.1
N743-1		0.62	0.1
N744-1		0.71	0.1
N745-1		6.94	0.5
PIN745-1		1.08	0.5
PIN747-1		0.68	0.1
PIN748-1		0.63	0.2
N749-1		1.64	0.1
PIN750-1		0.93	0.2
PIN751-1		1.05	0.1
PIN752-1	100	1.64	0.0
PIN753-1		1.49	0.3



ssed & Understood by me.

Date

invented by

Date

This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

BLACK BORDERS

IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

FADED TEXT OR DRAWING

BLURRED OR ILLEGIBLE TEXT OR DRAWING

SKEWED/SLANTED IMAGES

COLOR OR BLACK AND WHITE PHOTOGRAPHS

GRAY SCALE DOCUMENTS

LINES OR MARKS ON ORIGINAL DOCUMENT

REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

IMAGES ARE BEST AVAILABLE COPY.

☐ OTHER:

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.